

**Green-tight measure of Kato class and  
compact embedding theorem for symmetric  
Markov processes  
(joint work with Kazuhiro Kuwae)**

**Kaneharu Tsuchida  
(National Defense Academy)**

**Japanese-German Open Conference on Stochastic Analysis 2019  
at Fukuoka University**

**September 5, 2019**

- In this talk, we would like to discuss compact embeddings for symmetric Markov processes.
- Let  $(\mathcal{E}, \mathcal{F})$  be a Dirichlet form on  $L^2(E; m)$  associated with a  $m$ -symmetric Markov process  $X$ .
- For a suitable measure  $\mu$ , Stollmann-Voigt proved the following inequality: for  $\alpha > 0$

$$\int u^2 d\mu \leq \|R_\alpha \mu\|_\infty \mathcal{E}_\alpha(u, u), \quad u \in \mathcal{F}, \quad (1)$$

where  $R_\alpha$  is the  $\alpha$ -resolvent of  $X$  and

$\mathcal{E}_\alpha(u, u) = \mathcal{E}(u, u) + \alpha(u, u)$ . Moreover, if  $X$  is transient, (1) holds for  $\alpha = 0$  and  $u \in \mathcal{F}_e$ .

- Hence the embedding  $(\mathcal{F}, \mathcal{E}_1) \hookrightarrow L^2(\mu)$  (or  $(\mathcal{F}_e, \mathcal{E}) \hookrightarrow L^2(\mu)$  if  $X$  is transient) is continuous.

- Takeda introduced following conditions:

(I)  $X$  is **irreducible**:

If any Borel set  $B$  satisfies  $P_t 1_B u = 1_B P_t u$  for all  $u \in L^2(E; m)$  and  $t > 0$ , then  $m(B) = 0$  or  $m(B^c) = 0$  holds.

(RSF)  $X$  has the **resolvent strong Feller property**:

$R_\alpha(\mathcal{B}_b) \subset C_b$  for any  $\alpha > 0$ .

(Tightness)  $X$  has a **tightness property**:

For any  $\varepsilon > 0$ , there exists a compact set  $K(\subset E)$  such that

$$\|R_1(1_{K^c} m)\|_\infty < \varepsilon.$$

If  $X$  satisfies conditions (I), (RSF) and (Tightness),  $X$  is called **“class (T)”**.

## Theorem 1 (Takeda ('19))

Suppose that  $X$  is class (T).

- (1) The Markov semigroup is compact on  $L^2(E; m)$  and its every eigenfunction has a bounded continuous version.
- (2) The embedding  $(\mathcal{F}, \mathcal{E}_1) \hookrightarrow L^2(E; m)$  is compact.
- (3) If  $X$  is transient and  $\mu \in S_{CK_\infty}^1(X)$ , then the embedding  $(\mathcal{F}_e, \mathcal{E}) \hookrightarrow L^2(\mu)$  is compact.
- (4) There exists a bounded ground state uniquely up to sign, that is, the function  $\phi_0$  which attains the infimum:

$$\inf \left\{ \mathcal{E}(u, u) : u \in \mathcal{F}, \int_E u^2 dm = 1 \right\}.$$

Moreover,  $\phi_0$  can be taken to be strictly positive.

## Remark 2

- (1) In Theorem 1,  $(1) \iff (2)$ .
- (2) The statement (3) plays very important role to prove the large deviation for additive functionals.
- (3) For (3), Chen-T. proved by another method that this embedding is compact if  $X$  is pure jump symmetric Markov process which satisfies a mild condition for jump kernel without (I) and (RSF).

- In proofs of compactness, we notice that Takeda does not use (RSF) essentially.
- He use (RSF) in proving that  $m$  belongs to the class of Green-tight Kato measure in the sense of Chen (in notation  $S_{CK_\infty}^1(X^{(1)})$ ).  
( $X^{(1)}$  means the 1-subprocess of  $X$ ).
- We would like to clarify where these conditions are used, and generalize these results.

$E$  : a locally compact separable metric space

$m$  : a positive Random measure on  $E$  with full support.

$X = (\mathbb{P}_x, X_t)$  :  $m$ -symmetric special standard process on  $E$ .

$\{P_t, t \geq 0\}$ : the semigroup of  $X$ .

$(\mathcal{E}, \mathcal{F})$  : the quasi-regular Dirichlet form generated by  $X$ :

$$\mathcal{F} = \left\{ u \in L^2(m) : \lim_{t \downarrow 0} \frac{1}{t} ((I - P_t)u, u)_{L^2(m)} < \infty \right\}$$

$$\mathcal{E}(u, v) = \lim_{t \downarrow 0} \frac{1}{t} ((I - P_t)u, v)_{L^2(m)}, \quad u, v \in \mathcal{F}.$$

$(\mathcal{F}_e, \mathcal{E})$  : the extended Dirichlet space of  $(\mathcal{E}, \mathcal{F})$ .

$R_\alpha$  : the  $\alpha$ -resolvent of  $X$ .

$S^1(X)$  : the family of positive smooth measures in the strict sense under the absolute continuity condition (AC).

In this talk, we always assume that any measure belongs to  $S^1(X)$ .

Let  $P_t(x, dy)$  be the transition function of  $X$ , that is,

$$P_t(x, B) = \mathbb{P}_x(X_t \in B).$$

In the sequel, we use the following notations:

**(AC)** : for any  $t > 0$  and  $x \in E$ ,  $P_t(x, dy)$  is absolutely continuous with respect to  $m$ .

**(SF)** : for any  $t > 0$ ,  $P_t(\mathcal{B}_b(E)) \subset C_b(E)$ .

**(RSF)** : for any  $\alpha > 0$ ,  $R_\alpha(\mathcal{B}_b(E)) \subset C_b(E)$ .

It is known that

$$\text{(SF)} \implies \underbrace{\text{(RSF)}}_{\text{Takeda's results}} \implies \underbrace{\text{(AC)}}_{\text{Our results}} .$$



We define  $\alpha$ -potential of  $\nu$  by

$$R_\alpha \nu(x) = \mathbb{E}_x \left[ \int_0^\infty e^{-\alpha t} dA_t^\nu \right], \quad x \in E$$

where  $A_t^\nu$  is the PCAF associated to  $\nu \in S^1(X)$ .

### Definition 3 (Kato class)

- (1) Suppose that  $X$  is transient.  $\nu$  is said to be a **Green-bounded** ( $S_{D_0}(X)$ ) if  $\sup_{x \in E} R\nu(x) < \infty$ .
- (2)  $\nu$  is said to be a smooth measure of Kato class  $S_K^1(X)$  if

$$\lim_{\alpha \rightarrow \infty} \sup_{x \in E} R_\alpha \nu(x) = 0.$$

- (3) The local Kato class  $S_{LK}^1(X)$  is defined by

$$S_{LK} = \{\nu \in S^1(X) : 1_K \nu \in S_K^1(X) \text{ for any } K \text{ cpt.}\}.$$

## Definition 4 (Two kinds of Green-tight measure)

Let  $\nu \in S^1(X)$  and  $\alpha \geq 0$ . When  $\alpha = 0$ , we always assume the transience of  $X$ .

- (1) (Zhao)  $\nu \in S^1_{K_\infty}(X) \stackrel{\text{def.}}{\iff} \nu \in S^1_K(X)$  and for any  $\varepsilon > 0$  there exists a compact subset  $K = K(\varepsilon)$  of  $E$  such that

$$\sup_{x \in E} R_\alpha(1_{K^c} \nu)(x) < \varepsilon.$$

- (2) (Chen)  $\nu \in S^1_{CK_\infty}(X) \stackrel{\text{def.}}{\iff}$  for any  $\varepsilon > 0$  there exists a Borel subset  $K = K(\varepsilon)$  of  $E$  with  $\nu(K) < \infty$  and a constant  $\delta > 0$  such that for all  $\nu$ -measurable set  $B \subset K$  with  $\nu(B) < \delta$ ,

$$\sup_{x \in E} R_\alpha(1_{B \cup K^c} \nu)(x) < \varepsilon.$$

If  $\alpha > 0$ , we rewrite  $S_{K_\infty}^1(X)$  (resp.  $S_{CK_\infty}^1(X)$ ) with  $S_{K_\infty^+}^1(X)$  (resp.  $S_{CK_\infty^+}^1(X)$ ).

### Remark 5

- (1) **Definition 4(1):** Zhao originally introduced the class  $S_{K_\infty}^1(X)$  in considering the gaugeability for  $d$ -dim. absorbing Brownian motions ( $d \geq 3$ ) on bounded open domains.
- (2) **Definition 4(2):** However,  $S_{K_\infty}^1(X)$  is not enough to develop the gaugeability and subcriticality for symmetric Markov processes. To overcome some difficulty, Chen introduced the class  $S_{CK_\infty}^1(X)$ .
- (3) The Borel set  $K = K(\varepsilon)$  in Definition 4(2) can be taken to be a compact set by the inner regularity of  $m$ . Hence  $S_{CK_\infty^{(+)}}(X) \subset S_{K_\infty^{(+)}}(X)$ .

### Remark (continued)

- (4) Chen proved that  $S_{K_{\infty}^{(+)}}^1(X) = S_{CK_{\infty}^{(+)}}^1(X)$  under (SF). Later, Kim and Kuwae proved the coincidence under (RSF). Moreover, the equality holds under the ultracontractivity of  $X$ .
- (5) If  $\alpha > 0$ ,  $S_{K_{\infty}^{+}}^1(X)$  and  $S_{CK_{\infty}^{+}}^1(X)$  are independent of the choice of  $\alpha > 0$  by the resolvent equation.
- (6) Chen proved that  $(S_{CK_{\infty}}^1(X) \subset ) S_{CK_{\infty}^{+}}^1(X) \subset S_K^1(X)$ .
- (7) Clearly,  $S_{CK_{\infty}^{+}}^1(X) = S_{CK_{\infty}}^1(X^{(1)})$ .

- In the sequel, we only consider 0-order Green-tight measure by Remark 5(7).

## Theorem 6

Suppose that  $X$  satisfies (AC) and  $m \in S_{CK_\infty}^1(X^{(1)})$ . Then the  $L^2$ -semigroup  $P_t$  is a compact operator on  $L^2(E; m)$  and its every eigenfunction has a finely continuous Borel measurable bounded  $m$ -version. Moreover, if  $X$  satisfies (RSF), then every eigenfunction has a bounded continuous  $m$ -version.

## Theorem 7

Suppose that  $X$  satisfies (AC) and  $m \in S_{CK_\infty}^1(X^{(1)})$ . Then the embedding  $\mathcal{F} \hookrightarrow L^2(E; m)$  is compact.

## Theorem 8

Suppose that  $X$  is transient and it satisfies (AC). Let  $\nu \in S_{CK_\infty}^1(X)$ . Then  $(\mathcal{F}_e, \mathcal{E})$  is compactly embedded in  $L^2(E; \nu)$ .

Let  $\lambda_2$  be the bottom of the spectrum:

$$\lambda_2 := \inf \left\{ \mathcal{E}(f, f) : f \in \mathcal{F}, \int_E f^2 dm = 1 \right\}.$$

A function  $\phi_0$  on  $E$  is called a ground state of the  $L^2$ -generator for  $\mathcal{E}$  if  $\phi_0 \in \mathcal{F}$ ,  $\|\phi_0\|_2 = 1$  and  $\mathcal{E}(\phi_0, \phi_0) = \lambda_2$ .

## Theorem 9

Suppose that  $X$  satisfies (AC), (I) and  $m \in S_{CK_\infty}^1(X)$ . Then there exists a bounded ground state  $\phi_0$  uniquely up to sign. Moreover,  $\phi_0$  can be taken to be strictly positive on  $E$ .

### Theorem 10

Suppose that  $X$  is transient which possesses (RSF). Take  $\nu \in S_{D_0}(X)$  and assume  $\nu \notin S_{LK}(X)$ .

- (1) If  $\nu$  has the full quasi-support, then the time changed process  $(\check{X}, \nu)$  does not possess (RSF), but satisfies (AC).
- (2) There exists an  $\beta > 0$  such that the killed process  $X^{-\beta\nu}$  does not possess (RSF), but satisfies (AC).

Is there a measure  $\nu$  that satisfies this theorem?

## Example 1 (Brownian motion)

Let  $X$  be the  $d$ -dimensional BM on  $\mathbb{R}^d$  with  $d \geq 3$  and  $m$  the Lebesgue measure on  $\mathbb{R}^d$ .

Set  $x_n := (2^{-n}, 0, \dots, 0) \in \mathbb{R}^d$  and  $r_n = 8^{-n}$ . We set  $V_n(x) = 8^{2n} 1_{B_{r_n}(x_n)}(x)$  and  $V(x) := \sum_{n=2}^{\infty} V_n(x)$ . Then we find that  $Vm \in S_{D_0}(X) \setminus S_{LK}^1(X)$  by Aizenman-Simon ('82). Since  $X$  is transient, there exists a function  $g$  such that  $0 < g \leq 1$   $m$ -a.e. and  $Rg \in \mathcal{B}_b(E)$ . We put  $\nu = (V + g)m$ . Then we know that the time-changed processes  $\hat{X}^\nu$  associated with  $\nu$  and the killed process  $X^{-\beta\nu}$  for some  $\beta > 0$  do not possess (RSF) by Theorem 10, but satisfy (AC).



## Example 2 (stable process)

Take  $\alpha \in (0, 2)$  and  $m \geq 0$ . Let  $X = (\Omega, X_t, \mathbb{P}_x)$  be a Lévy process on  $\mathbb{R}^d$  with

$$\mathbb{E}_0[e^{i\langle \xi, X_t \rangle}] = \exp\left(-t(|\xi|^2 + m^{2/\alpha})^{\alpha/2} - m\right)$$

If  $m > 0$ , it is called the relativistic  $\alpha$ -stable process with mass  $m$ . We assume the transience of  $X$ , i.e.  $d \geq 3$  with  $m > 0$ , or  $d > \alpha$  with  $m = 0$ . Let  $x_n$  and  $r_n$  be the point and constant as in Example 1. We fix  $G := B_1(0)$ . We set  $V_n(x) = 8^{\alpha n} 1_{B_{r_n}(x_n)}(x)$  and  $V(x) := \sum_{n=2}^{\infty} V_n(x)$ . Then  $Vm \in S_{D_0}(X) \setminus S_{LK}^1(X)$ . Putting  $\nu = (V + g)m$ , we know that the time-changed process  $\check{X}^\nu$  and killed process  $X^{-\beta\nu}$  for some  $\beta > 0$  do not possess (RSF) by Theorem , but satisfy (AC).

**Thank you for your attention !!**